

Research



**Cite this article:** Schumann I, Kenny N, Hui J, Hering L, Mayer G. 2018 Halloween genes in panarthropods and the evolution of the early moulting pathway in Ecdysozoa. *R. Soc. open sci.* 5: 180888.  
<http://dx.doi.org/10.1098/rsos.180888>

Received: 8 June 2018

Accepted: 17 August 2018

**Subject Category:**

Biology (whole organism)

**Subject Areas:**

molecular biology/evolution/developmental biology

**Keywords:**

ecdysis, ecdysone, 20-hydroxyecdysone, Halloween genes, Onychophora, Tardigrada

**Author for correspondence:**

Georg Mayer

e-mail: [georg.mayer@uni-kassel.de](mailto:georg.mayer@uni-kassel.de)

<sup>†</sup>These authors contributed equally to this study.

<sup>‡</sup>Present address: Life Sciences, The Natural History Museum, Cromwell Road, Kensington, London SW7 5BD, UK.

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.4217987>.

# Halloween genes in panarthropods and the evolution of the early moulting pathway in Ecdysozoa

Isabell Schumann<sup>1,2,†</sup>, Nathan Kenny<sup>3,‡</sup>, Jerome Hui<sup>3</sup>, Lars Hering<sup>1,†</sup> and Georg Mayer<sup>1</sup>


<sup>1</sup>Department of Zoology, Institute of Biology, University of Kassel, Kassel, Germany

<sup>2</sup>Molecular Evolution and Animal Systematics, Institute of Biology, University of Leipzig, Leipzig, Germany

<sup>3</sup>School of Life Sciences, Simon F.S. Li Marine Science Laboratory, Center of Soybean Research, State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong, People's Republic of China

IS, 0000-0001-8354-2523; NK, 0000-0003-4816-4103; JH, 0000-0003-1355-8495

Moulting is a characteristic feature of Ecdysozoa—the clade of moulting animals that includes the hyperdiverse arthropods and less speciose groups, such as onychophorans, tardigrades and nematodes. Moulting has been best analysed in arthropods, specifically in insects and crustaceans, in which a complex neuroendocrine system acts at the genomic level and initiates the transcription of genes responsible for moulting. The key moulting hormones, ecdysone and 20-hydroxyecdysone, are subsequently synthesized from cholesterol ingested with food. Their biosynthesis is regulated by the Rieske-domain protein Neverland and cytochrome P450 enzymes encoded by the so-called ‘Halloween’ genes. Ecdysone is then released into the haemolymph and modified into 20-hydroxyecdysone, which binds to the nuclear receptor EcR/USP and initiates transcription of the Early genes. As little is known about the moulting pathway of other ecdysozoans, we examined the occurrence of genes involved in ecdysteroid biosynthesis and the early moulting cascade across ecdysozoan subgroups. Genomic and transcriptomic searches revealed no Halloween genes in cycloneuralians, whereas only *shadow* (*CYP315A1*) is present in onychophorans and tardigrades, suggesting that the Halloween genes evolved stepwise in panarthropods. These findings imply that the genes which were responsible for the ecdysteroid biosynthesis in the last common ancestor of Ecdysozoa are currently unknown.




**Figure 1.** Identified occurrence of 20-hydroxyecdysone in protostomes. Note that 20-hydroxyecdysone (indicated by red dots) is not only found in ecdysozoans but also in other protostomes, including molluscs, annelids and platyhelminths. Phylogeny of protostomes modified from [1,2]. Note that cycloneurians might not be monophyletic [1]. Animal silhouettes, except for those for Onychophora and Tardigrada, courtesy of PhyloPic ([www.phylopic.org](http://www.phylopic.org)).

## 1. Introduction

Ecdysozoa is the sister group of Lophotrochozoa within protostomes and includes cycloneurians (nematodes, priapulids, kinorhynchs and allies), and panarthropods (tardigrades, onychophorans and arthropods; figure 1). Although the phylogenetic relationships of ecdysozoans remain contentious (in particular, the validity of Cycloneuralia is under debate), the monophyly of the entire clade is supported by most molecular analyses and the process of moulting or ecdysis, which is considered an autapomorphy of this clade [1,3–6]. Ecdysis is essential for growth in these animals, as their body is typically covered with an inelastic cuticle or exoskeleton, which has to be shed and replaced periodically by larger covering. During this complex and strictly coordinated neuroendocrine process, precise timing of gene expression and physiological responses is essential for building the new cuticle or exoskeleton [7,8]. Hormones such as ecdysteroids have been demonstrated to be responsible for the control and regulation of moulting in arthropods [5,9]. Specifically, ecdysone (E) and 20-hydroxyecdysone (20E) have been identified as key players of ecdysis in insects [10] and crustaceans (crustecdysone *sensu* Hampshire & Horn [11]; figure 2*a,b*). These hormones are synthesized from the precursor sterol cholesterol, which is ingested with food, transported to a secretory tissue and converted subsequently into the final moulting hormone 20E [14] (figure 2*b*).

In well-studied insects, such as the fruit fly *Drosophila melanogaster* and the tobacco hornworm *Manduca sexta*, the initial step in the biosynthesis of ecdysteroids is the dehydrogenation of cholesterol to 7-dehydrocholesterol, which takes place in the prothoracic gland and is catalysed by the Rieske oxygenase Neverland (Nvd) [15–17] (figure 2*b*). The subsequent hydroxylation reactions at different carbon atoms are catalysed by cytochrome P450 monooxygenases encoded by the Halloween genes *spook* (*spo*, CYP307A1), *phantom* (*phm*, CYP306A1), *disembodied* (*dib*, CYP302A1) and *shadow* (*sad*, CYP315A1). In *D. melanogaster*, two additional Halloween genes—*spookier* (*spok*, CYP307A2) and *spookiest* (CYP307B1)—have been identified with a putative hydroxylation function as part of the ‘Black box’ in the ecdysteroid pathway [18–20] (figure 2*b*). These multiple hydroxylation reactions in the secretory tissue produce ecdysone. This is released into the haemolymph and modified further to 20E by a monooxygenase encoded by the Halloween gene *shade* (*shd*, CYP314A1) in a species-specific target area such as the haemolymph, fat body or midgut [21,22]. The active moulting hormone 20E



**Figure 2.** Hormonal titre and early moulting pathway in the fruit fly *Drosophila melanogaster*. Modified from [12,13]. (a) Changes of 20-hydroxyecdysone titre during development. Note that the titre increases shortly before each moult and metamorphosis. (b) Ecdysteroid biosynthesis and early moulting pathway in a 3rd instar larva (time point indicated by arrow in (a)). The numbers 1–5 indicate major steps of biosynthesis and binding to ecdysteroid receptor. Ingestion of dietary cholesterol is followed by ecdysone biosynthesis in secretory tissue (prothoracic gland), which is controlled by enzymes encoded by the Rieske-domain gene *neverland* (*nvd*), and the Halloween genes *spook* (*spo*), *spookier* (*spok*), *phantom* (*phm*), *disembodied* (*dib*) and *shadow* (*sad*). Ecdysone is then released into the haemolymph and oxidized to 20-hydroxyecdysone by an oxidase encoded by *shade* (*shd*). Binding of 20-hydroxyecdysone to the receptor complex EcR/USP in target tissue (epidermis) activates transcription of the Early genes (*E74*, *E75*, *E78*, *HR3*, *HR4* and  $\beta$ Ftz-F1), which is followed by cuticle biosynthesis. 2-dE, 2-deoxyecdysone; 2,22-dE, ketotriol; 2,22,25-dE, ketodiol; 7-dC, 7-dehydrocholesterol; 20E, 20-hydroxyecdysone; C, cholesterol; E, ecdysone; *E74*, ecdysone-inducible gene 74; *E75*, ecdysone-inducible gene 75; *E78*, ecdysone-inducible gene 78; *HR3*, hormone receptor 3; *HR4*, hormone receptor 4;  $\beta$ Ftz-F1, beta Fushi-tarazu transcription factor 1; EcR, ecdysone receptor; ERE, ecdysone response element; USP, Ultraspiracle.

then binds to a high-affinity nuclear receptor complex that consists of two dimerization partners, Ecdysone Receptor (EcR) and the retinoid X receptor homologue Ultraspiracle (USP) [23–25]. By attaching to an ecdysone response element of the DNA, the 20E/EcR/USP complex initiates a hierarchical transcription cascade of the Early genes (at least six in *D. melanogaster*, including *E74*, *E75*, *E78*, *HR3*, *HR4* and  $\beta$ Ftz-F1), which leads to the final biosynthesis of the new cuticle by the epidermis regulated by enzymes (e.g. chitinases and collagenases) encoded by the Late response genes [26–29] (figure 2b).

The molecular control of moulting is well understood in insects and crustaceans (e.g. [30–32]). However, little is known about this process in the remaining ecdysozoan subgroups, including the closest arthropod relatives, the onychophorans (velvet worms) and tardigrades (water bears) [1,12,33]. While in the onychophoran *Euperipatoides leuckartii* both key moulting hormones, E and 20E, have been identified using radioimmunoassay and high-performance liquid chromatography [34], corresponding data are unavailable from tardigrades, probably due to their minute body size

**Table 1.** Ecdysozoan species, the genomes and/or transcriptomes of which were searched for homologues of genes of the moulting pathway.

taxon	species	genome	transcriptome	references
Scalidophora, Priapulida	<i>Priopus caudatus</i>	✓		AXZU00000000.2 <a href="http://genome.wustl.edu/">http://genome.wustl.edu/</a>
Nematoda, Rhabditida	<i>Caenorhabditis elegans</i>	✓		[47]
Tardigrada, Eutardigrada	<i>Hypsibius exemplaris</i>	✓	✓	[48] [49]
	<i>Ramazzottius varieornatus</i>	✓		[50]
Onychophora, Peripatopsidae	<i>Euperipatoides rowelli</i>	✓	✓	<a href="ftp://ftp.hgsc.bcm.edu/15K-pilot/Velvet_worm/">ftp://ftp.hgsc.bcm.edu/15K-pilot/Velvet_worm/</a> [51]
Arthropoda, Chelicerata	<i>Ixodes scapularis</i>	✓		[52]
	<i>Tetranychus urticae</i>	✓		[53]
	<i>Stegodyphus mimisarum</i>	✓		[54]
Arthropoda, Myriapoda	<i>Strigamia maritima</i>	✓		[55]

(figure 1). To our knowledge, the occurrence of ecdysteroids has not been investigated in cycloneurialian taxa other than nematodes. Although ecdysteroids have been detected in the parasitic nematodes *Diofilaria immitis* and *Ascaris suum* [35,36], neither E nor 20E have been identified in free-living nematodes, including the ‘model’ species *Caenorhabditis elegans* [37,38]. This would suggest a different pathway for the regulation of moulting in the free-living species of nematodes at least. Dafachronic acid, another sterol-derived hormone, has, for example, been identified as a moulting hormone in *C. elegans* [39].

Interestingly, the ecdysteroids E and 20E have also been detected outside the ecdysozoan clade, for example, in the cestode *Moniezia expansa*, the molluscs *Lymnaea stagnalis* and *Helix pomatia*, and the leech *Hirudo medicinalis* [40–44] (figure 1). Together with the observation that leeches periodically shed their cuticle [43,45,46], this raises the question of whether or not the ecdysteroid pathway is conserved among protostomes, and whether ecdysone-induced moulting predates the origin of Ecdysozoa.

To clarify this question, we explored the distribution of the major gene components of the early moulting pathway in insects across the Bilateria. In particular, we focused on the occurrence of the Halloween genes among ecdysozoans by including the sequenced genomes and transcriptomes of a priapulid, two tardigrade and one onychophoran species in our analyses (table 1). After identifying the relevant sequences, we performed phylogenetic analyses of the three major families of moulting genes, including those encoding a Rieske-domain protein, cytochrome P450 monooxygenases and nuclear hormone receptors. Our results contribute to a better understanding of the evolution of genes of the early moulting pathway in panarthropods. They also reveal substantial gaps in our knowledge of the ancestral moulting system of ecdysozoans.

## 2. Material and methods

### 2.1. Specimens, library preparation, sequencing and transcriptome assembly

Specimens of the onychophoran species *Euperipatoides rowelli* Reid, 1996 (Peripatopsidae) were collected from decaying logs in the Tallaganda State Forest (New South Wales, Australia, 35°26' S, 149°33' E, 954 m above sea level) in October 2011. The animals were kept in plastic jars with perforated lids at 18°C or as

described previously [56,57] and fed with crickets (*Acheta domesticus*) every four weeks. Specimens of the eutardigrade species *Hypsibius exemplaris* Gąsiorek *et al.* [58] (Parachela, Hypsibiidae) were obtained commercially from Sciento (Manchester, UK) and kept in plastic Petri dishes filled with mineral water (Volvic, Danone Waters Deutschland GmbH, Frankfurt am Main, Germany) at 21°C and fed with unicellular algae (*Chlorococcum* sp.) as described previously [59–61]. This tardigrade species was commonly referred to as '*Hypsibius dujardini* (Doyère, 1840)' in the literature (e.g. [48,49,62]), but Gąsiorek *et al.* [58] have described it as *Hypsibius exemplaris*. Library preparation, sequencing and assembly of transcriptomes for *E. rowelli* and *H. exemplaris* were performed as described previously [49,51].

## 2.2. Identification of genes and genome screening

To identify the sequences of putative genes from the early moulting pathway of various ecdysozoans (including *neverland*, Halloween genes, receptor genes, Early genes, and *CYP18A1*) (table 1), tBLASTn/BLASTp searches [63] of the NCBI GenBank database were conducted using an initial score cut-off of 80 and *E*-value of  $1 \times 10^{-6}$ . Published sequences of the Halloween genes from different arthropods, including *Limulus polyphemus*, *Daphnia pulex* and *D. melanogaster*, were used as queries (see electronic supplementary material, file F1 and figures S1–S3 for accession numbers or references for all used sequences). When the target gene was not found by initial searches, further searches at a more permissive score of 60 and higher *E*-value cut-offs were performed. Alongside these searches, the available genomes of different ecdysozoan species, including *Priapululus caudatus*, *C. elegans*, *H. exemplaris*, *Ramazzottius varieornatus*, *E. rowelli*, *Ixodes scapularis*, *Tetranychus urticae*, *Stegodyphus mimisarum* and *Strigamia maritima* were screened for candidate genes using Hmmer or BLAST searches either directly on the Ensembl database [64] or on their respective webservers (table 1). Reciprocal tBLASTn/BLASTP searches of candidate gene sequences on UniProt and NCBI databases were performed for initial confirmation of gene identity from the examined genomes (table 1). Candidate sequences from the onychophoran *E. rowelli* and the tardigrade *H. exemplaris* were translated using the standard metazoan coding table with the online sequence translation tool EMBOSS Transeq ([65], [http://www.ebi.ac.uk/Tools/st/emboss\\_transeq](http://www.ebi.ac.uk/Tools/st/emboss_transeq)) and CLC Sequence viewer (Qiagen, 2017). The domain structures of all sequences (Rieske-domain for *neverland*, P450 domain for the Halloween genes and *CYP18A1*, Zn-Finger motif and hormone receptor for the receptor genes and the Early genes, ETS-domain for *E74*) were analysed using the Pfam database 31.0 [66] and SMART [67]. We used the corresponding sequences of the ETS transcription factor family to clarify the orthology of the identified *E74* sequences (*ETS*, *TCF*, *PEA3* and *ERG sensu* Sharrocks *et al.* [68]).

## 2.3. Sequence alignment and phylogenetic analyses

Amino acid sequences of putative genes of the early moulting pathway (59 for *neverland*, 238 for the Halloween genes and *CYP18A1*, 173 for the nuclear hormone receptors and 58 for the ETS transcription factor family genes) were used for phylogenetic analyses. We generated dataset and domain sequence alignments for each gene family (257 amino acids in length for *neverland*, 1006 for the Halloween genes and *CYP18A1*, 665 for the nuclear hormone receptors and 83 for the ETS transcription factor family genes) using the online tool MAFFT version 7 [69] and the L-INS-I strategy (see electronic supplementary material, file F2 for all alignments). The Pthread-Version of RAxML v. 8.2.X [70] was used to infer the best maximum-likelihood tree to reveal the phylogenetic position of each candidate sequence by using a dataset-specific GTR substitution matrix (-m PROTGAMMAGTR). For each run, the best tree was obtained from 10 independent inferences. Bootstrap support values for all trees were calculated using the rapid bootstrapping algorithm implemented in RAxML from 1000 pseudoreplicates. Trees were visualized with iTol v. 4.0.3 [71] and edited with Illustrator CS5 (Adobe Systems, San Jose, CA, USA).

# 3. Results

## 3.1. Homologues of the Rieske-domain gene *neverland*

Our genomic/transcriptomic searches and phylogenetic analyses revealed homologues of *neverland* (*nvd*) in numerous bilaterians (table 2; see electronic supplementary material, figure S1 for phylogenetic analyses). We identified orthologues of *nvd* in all analysed arthropod species, including the chelicerate *I. scapularis*, the

**Table 2.** Identified homologs (✓) of candidate genes involved in the moulting process across major ecdysozoan taxa. Note that genomic or deep transcriptomic data are unavailable for Loricifera, Kinorhyncha, and Nematomorpha. Abbreviations: *βFtz-F1*, *beta fushi tarazu transcription factor 1*; *CYP18A1*, *cytochrome P450-18A1*; *dib*, *disembodied*; *Ecr*, *ecdysone receptor*; *HR3*, *hormone receptor 3*; *HR4*, *hormone receptor 4*; *nvd*, *neverland*; *phm*, *phantom*; *sad*, *shadow*; *shd*, *shade*; *spo*, *spook*; *spookier*, *USP/RXR*, *ultraspiracle/retinoid X receptor*.

Taxon	Gene									
	Priapulida	Nematoda	Tardigrada	Onychophora	Chelicerata	Myriapoda	'Crustacea'	Hexapoda		
<b>Rieske-domain gene</b>										
<i>nvd</i>	✗	✓	✗	✗	✓	✓	✓	✓		
<b>Halloween genes</b>										
<i>sad</i> <sup>1</sup>	✗	✗	✓	✓	✓	✓	✓	✓		
<i>spo</i>	✗	✗	✗	✗	✓	✓	✓	✓		
<i>dib</i>	✗	✗	✗	✗	✓	✓	✓	✓		
<i>shd</i>	✗	✗	✗	✗	✓	✓	✓	✓		
<i>phm</i> <sup>2</sup>	✗	✗	✗	✗	✗	✓	✓	✓		
<i>spookiest</i> <sup>3</sup>	✗	✗	✗	✗	✗	✓	✓	✓		
<i>spok</i> <sup>4</sup>	✗	✗	✗	✗	✗	✗	✗	✗		
<b>Receptor genes</b>										
<i>Ecr</i>	✓	✓	✓	✓	✓	✓	✓	✓		
<i>USP/RXR</i>	✓	✓	✓	✓	✓	✓	✓	✓		
<b>Early genes</b>										
<i>E74</i>	✓	✓	✓	✓	✓	✓	✓	✓		
<i>E75</i>	✓	✓	✓	✓	✓	✓	✓	✓		
<i>E78</i>	✓	✓	✓	✓	✓	✓	✓	✓		
<i>HR3</i>	✓	✓	✓	✓	✓	✓	✓	✓		
<i>HR4</i>	✓	✓	✓	✓	✓	✓	✓	✓		
<i>βFtz-F1</i>	✓	✓	✓	✓	✓	✓	✓	✓		
<b>Degradation gene</b>										
<i>CYP18A1</i> <sup>5</sup>	✗	✗	✓	✓	✓	✓	✓	✓		

<sup>1</sup> Three copies of *sad* are present in the onychophoran *Euperipatoides rovellii* (Peripatopsidae).

<sup>2</sup> Homologues of *phm* are missing in the genomes of the chelicerates *Stegodyphus mimosarum*, *Ixodes scapularis* and *Tetranychus urticae*.

<sup>3</sup> *spookiest* might have been lost in the cladocerans *Daphnia pulex* and *D. magna*, as its homologues are missing in their genomes.

<sup>4</sup> Homologues of *spook* seem to be present only in drosophilids.

<sup>5</sup> *CYP18A1* is responsible for the degradation of 20-hydroxyecdysone in insects.

myriapod *S. maritima*, the crustacean *D. pulex* and the insect *D. melanogaster*. While genomic and transcriptomic data from Kinorhyncha, Loricifera and Nematomorpha are still absent from available databases, we confirmed the previously identified homologue of *nvd*, *daf-36* (cf. [17]), in the nematode *C. elegans*. By contrast, we found no *nvd* homologues in genomic and/or transcriptomic data from the onychophoran *E. rowelli*, the tardigrades *H. exemplaris* and *R. varieornatus*, and the priapulid *P. caudatus* (table 2; see electronic supplementary material, figure S1 for phylogenetic analyses).

### 3.2. Cytochrome P450 family genes

To clarify the orthology of the Halloween genes, we analysed the phylogeny of selected cytochrome P450 family genes (figure 3). Our screening of genomic and transcriptomic data from the onychophoran *E. rowelli* and the tardigrades *H. exemplaris* and *R. varieornatus* revealed only orthologues of the Halloween gene *shadow* (*sad*, *CYP315A1*) in these three panarthropod species (figure 3 and table 2). While the two tardigrade species show single homologues of *sad*, we identified three transcripts of *sad* in the onychophoran *E. rowelli* (figure 3). The tardigrade and onychophoran homologues form a clade together with the corresponding sequences from arthropods (figure 3). Besides the homologues of *sad*, our results revealed four additional Halloween genes (*spo*, *dib*, *shd* and *phm*) in the myriapods *S. maritima* and *Chamberlinius hualienensis*, the crustaceans *D. pulex* and *Daphnia magna*, and several insect species, whereas *phm* is missing in the chelicerates *S. mimosarum*, *I. scapularis* and *T. urticae* (table 2; see electronic supplementary material, figure S2 for uncondensed phylogenetic analyses). In contrast to onychophorans, tardigrades and arthropods, the genomic data from nematodes and the priapulid *P. caudatus* exhibit no Halloween genes.

Our searches further revealed putative orthologues of the cytochrome P450 gene *CYP18A1* at least in all major subgroups of panarthropods studied (figure 3 and table 2). Interestingly, the *daf-9* sequences from the nematodes *C. elegans*, *Brugia malayi* and *Strongyloides ratti* (indicated by an orange asterisk in figure 3) fall into an assemblage of other nematode cytochrome P450 family genes, which cluster as the sister group to a clade comprising *CYP18A1*, *phm* and other P450 sequences (figure 3). We did not find homologues of *CYP18A1* in the priapulid *P. caudatus* (figure 3 and table 2).


### 3.3. Nuclear receptor and ETS transcription factor genes

We further searched the available databases for homologues of the nuclear hormone receptor genes *EcR*, *ultraspiracle/retinoid X receptor* (*USP/RXR*), and the Early genes *E75*, *E78*, *HR3*, *HR4* and *βFtz-F1* (figure 4 and table 2). We used *knirps* and related sequences, which are part of the nuclear hormone receptor family, from several arthropods as an outgroup. Besides representative ecdysozoan sequences, we included *EcR* and *USP/RXR* homologues from the annelid *Capitella teleta*, the brachiopod *Lingula anatina* and the vertebrates *Gallus gallus* and *Homo sapiens* in our sampling. We identified homologues of *EcR* and *USP/RXR*, which encode the two dimerization partners of the moulting receptor complex in arthropods, in all studied ecdysozoan subgroups (figure 4 and table 2). Interestingly, we found no orthologues of *EcR* and *USP* in the genome of the free-living nematode *C. elegans*, although the corresponding sequences are present in the parasitic nematodes *Trichinella spiralis*, *Dirofilaria immitis* and *Brugia malayi*. Among the Early genes, which also belong to the nuclear hormone receptor family, we identified *E75*, *E78*, *HR3*, *HR4* and *βFtz-F1* in all studied ecdysozoans (figure 4 and table 2). The same holds true for the homologues of the ecdysone-inducible Early gene, *E74*, which belongs to the ETS transcription factor family (figure 5 and table 2).

## 4. Discussion


### 4.1. Loss of *neverland* homologues in Tardigrada, Onychophora and Priapulida

The Rieske-domain gene *nvd* encodes an oxygenase, which in insects acts upstream of the ecdysteroid biosynthesis pathway and dehydrogenates cholesterol to 7-dehydrocholesterol [15,17] (cf. figure 2*b*). However, Nvd is also known to play a more general role in the transport and metabolism of cholesterol [72]. Based on our results, we propose that a homologue of *nvd* might have existed in the last common ancestor of bilaterians, in which it might have played an ancient role in the metabolism of steroids, as these hormones are essential for development, growth and homeostasis in both protostomes and deuterostomes [17]. Surprisingly, we did not find homologues of *nvd* in the



**Figure 3.** Phylogenetic relationship of the cytochrome P450 family genes. The tree is based on a maximum-likelihood analysis of 238 P450 domain sequences of the mitochondrial (Mito) and cytochrome P450 family 2 (CYP2) from different ecdysozoans (see electronic supplementary material, file F1 for identified sequences and accession numbers, and figure S2 for uncondensed tree). Numbers at nodes indicate bootstrap support values greater than 50% obtained from 1000 pseudoreplicates. Note that the homologues of the Halloween genes *disembodied* (*dib*), *phantom* (*phm*), *shade* (*shd*) and *spook* (*spo*) are missing in onychophorans and tardigrades. Single homologues of *CYP18A1* are present in the onychophoran *Euperipatoides rowelli* (highlighted in brown) and the tardigrades *Hypsibius exemplaris* and *Ramazzottius varieornatus* (both highlighted in blue). Three copies of *shadow* (*sad*, *CYP15A1*) are present in the onychophoran *E. rowelli* (highlighted in brown) and one copy in the tardigrades *H. exemplaris* and *R. varieornatus* (highlighted in blue). Note further that the nematode *daf-9* sequences occur within a clade of other nematode cytochrome P450 family genes, indicating that *daf-9* might not be an orthologue of *CYP18A1* (orange asterisk).





**Figure 4.** Phylogenetic relationship of the nuclear hormone receptor genes. The tree is based on a maximum-likelihood analysis of 173 sequences from different bilaterian species (see electronic supplementary material; file F1 for accession numbers and electronic supplementary material, figure S3 for uncondensed tree). Numbers at nodes indicate bootstrap support values greater than 50% obtained from 1000 pseudoreplicates. Note that the homologues of *ecdysone receptor* (*EcR*), *beta fushi tarazu transcription factor 1* (*Bftz-F1*), *hormone receptor 3* (*HR3*), *hormone receptor 4* (*HR4*), *ecdysone-inducible genes E75* and *E78*, and *ultraspiracle/retinoid X receptor* (*USP/RXR*) are present in the onychophoran *Euperipatoides rowelli* (highlighted in brown) and the tardigrades *Hypsibius exemplaris* and *Ramazzottius varieornatus* (both highlighted in blue).

onychophoran *E. rowelli*, the tardigrades *H. exemplaris* and *R. varieornatus* and the priapulid *P. caudatus*, suggesting independent losses of this gene in the corresponding lineages. Owing to these losses, the enzyme responsible for the transformation of cholesterol to 7-dehydrocholesterol in these three animal groups is currently unknown.

#### 4.2. Stepwise evolution of Halloween genes in Panarthropoda

The Halloween genes encode hydroxylases that, at least in crustaceans and insects, convert 7-dehydrocholesterol into ecdysone and 20-hydroxyecdysone [73–75]. Mutations of the Halloween genes in the fruit fly *D. melanogaster* result in ecdysteroid deficiency and lead to cuticle deformations and


embryonic lethality [76,77]. Despite their essential role in the fruit fly, our results revealed no homologues of Halloween genes in cycloneuralians, suggesting that these animals might use an alternative pathway for the biosynthesis of ecdysteroids, or they might even use another type of moulting hormones. The free-living nematode *C. elegans* indeed seems to use dafachronic acid instead of 20E as the main moulting hormone [38,78], whereas the parasitic nematodes *A. suum* and *B. malayi* do show a molecular response to ecdysone [79–81]. Hence, the question arises as to whether the dafachronic acid pathway or the ecdysteroid pathway was responsible for the biosynthesis of moulting hormones in the last common ancestor of Cycloneuralia, provided this group is monophyletic [1]. Genome sequencing and, in particular, experimental studies in additional cycloneuralian taxa, such as priapulids, kinorhynchs, loriciferans and nematomorphs, might help to clarify this question.

Our results further show that the repertoire of Halloween genes varies considerably among arthropods. For example, the canonical 20E biosynthesis pathway of *D. melanogaster* exhibits six Halloween genes: *spo*, *spok*, *phm*, *dib*, *sad* and *shd* (cf. figure 2b), whereas most other panarthropods possess distinct sets of Halloween genes. While *spookiest* is missing in *Drosophila* species, this gene does occur in other insects and some crustaceans, suggesting that *spookiest* was present in the last common ancestor of Pancrustacea but was lost in some lineages, including the drosophilids and the cladoceran crustacean *Daphnia* (figure 6). Moreover, our phylogenetic analyses revealed that *spok* is a drosophilid in-paralogue of *spo*, thus confirming that *spok* might have evolved by gene duplication in the drosophilid lineage [18,75]. Although homologues of *spo*, *phm*, *dib*, *sad* and *shd* are present in myriapods, crustaceans and insects, *phm* is apparently missing in chelicerates (as speculated in Qu *et al.* [82]). The lack of this gene implies that *phm* was either lost in chelicerates or evolved in mandibulates (figure 6). Finally, *sad* is the only Halloween gene we identified in the onychophoran *E. rowelli* and the tardigrades *H. exemplaris* and *R. varieornatus*. While single orthologues of *sad* are present in tardigrades, the onychophoran transcriptome and genome databases exhibit three copies of this gene, suggesting that there might have been two duplication events in the onychophoran lineage or an onychophoran subclade.

In summary, based on our findings we propose that the most plausible scenario is a stepwise evolution of Halloween genes in panarthropods (figure 6). While only *sad* was present in the last common ancestor of Panarthropoda—a condition which has been retained in extant onychophorans and tardigrades—the genes *spo*, *dib* and *shd* might have arisen in the arthropod lineage, followed by the evolution of *phm* in mandibulates. The Halloween gene *spookiest* might have originated in the pancrustacean lineage, while *spok* evolved in drosophilids. These findings suggest that the biosynthesis pathway of ecdysteroids in tardigrades, onychophorans and chelicerates must be different from the canonical pathway of *D. melanogaster*. Owing to the lack of *phm*, the conversion of 2,22,25-dE to 2,22-dE (cf. figure 2b) might be accomplished by another enzyme in chelicerates, which is unknown. Alternatively, the lack of *phm* in these animals might be due to the use of ponasterone A (25-deoxy-20E) instead of 20E as a moulting hormone [82]. The situation is even less clear in onychophorans and tardigrades. Even if their only Halloween gene *sad* does participate in the conversion of 2-dE to E (cf. figure 2b)—a function which still has to be demonstrated experimentally—the remaining members of the ecdysteroid pathway in these animals are unknown. Hence, virtually nothing is known about the components of the ecdysteroid pathway and the molecular mechanisms of moulting in the last common ancestor of Panarthropoda.

### 4.3. Evolution of the EcR/USP complex and the Early genes


Another important process in the ecdysteroid pathway of insects is binding of 20E to the heterodimeric receptor complex EcR/USP, which activates the hierarchical transcription cascade of the Early genes [25–27] (figure 2b). The individual subunits of this receptor complex, which is also essential for reproduction and embryogenesis [83–85], are encoded by the *EcR* and *USP/RXR* genes. While *EcR* encodes three isoforms of the EcR subunit in *D. melanogaster* and two in the tobacco hornworm *M. sexta* [86,87], our analyses revealed single transcripts of *EcR* in the onychophoran *E. rowelli*, and the tardigrades *H. exemplaris* and *R. varieornatus*. We further identified single homologues of *USP/RXR*—the gene encoding the dimerization partner of EcR—in the onychophoran and the two tardigrade species. The occurrence of 20E and homologues of *EcR* and *USP/RXR* in several ecdysozoan taxa, including priapulids and the parasitic nematodes, suggests that the transcription of the Early genes might have been triggered by binding of 20E (or a related molecule) to the EcR/USP complex in the last common ancestor of Ecdysozoa. This ancient role of EcR/USP in steroid binding might have been inherited from



**Figure 5.** Phylogenetic relationship of ETS family genes including E74. The tree is based on a maximum-likelihood analysis of 58 sequences from different bilaterians (see electronic supplementary material, file F1 for identified sequences and accession numbers). Note the presence of E74 homologues in the onychophoran *Euperipatoides rowelli* (highlighted in brown) and the tardigrades *Hypsibius exemplaris* and *Ramazzottius varieornatus* (both highlighted in blue).

the last common ancestor of protostomes, as genes encoding both receptor subunits are also found in various other protostomes outside the ecdysozoan clade.

After steroid binding, the 20E/EcR/USP complex binds to an ecdysone response element in the promoter region and initiates the expression of the Early genes that encode transcription factors controlling the time and specificity of moulting in insects [29,88–90]. Our genomic and transcriptomic analyses revealed homologues of six Early genes in the onychophoran *E. rowelli*, and the tardigrades *H. exemplaris* and *R. varieornatus*: five of which were classified as the nuclear hormone receptor genes (*E75*, *E78*, *HR3*, *HR4* and  $\beta$ Ftz-F1), and one belonging to the ETS family of transcription factors (*E74*). Although the function of these genes has been well analysed in insects [90–93], only little is known about their potential role in moulting in other ecdysozoans. Studies on the nematode *C. elegans* have demonstrated that beyond their roles in development and egg-laying, at least some of the Early genes are involved in moulting [94,95]. Together with our finding of a complete set of Early genes in all studied ecdysozoan subgroups and other protostomes, such as the brachiopod *L. anatina* and the annelid *C. teleta*, this suggests that the Early genes might have been present in the last common ancestor of protostomes and were recruited for moulting in Ecdysozoa, irrespective of whether their transcription was initiated by 20E or another steroid hormone.



**Figure 6.** Scenario describing stepwise evolution of genes of the early moulting pathway in Ecdysozoa. Note that the last common ancestor of Panarthropoda possessed only the Halloween gene *shadow* (*sad*) and that the remaining Halloween genes, including *spook* (*spo*), *disembodied* (*dib*), *shade* (*shd*), *phantom* (*phm*), *spookiest* and *spookier* (*spok*), evolved stepwise in the arthropods.

#### 4.4. Evolution of the ecdysteroid-inactivating enzyme cytochrome P450-18A1 in panarthropods

In addition to ecdysone oxidase, the cytochrome P450 protein CYP18A1 has been characterized as an ecdysteroid-inactivating enzyme, which plays an important role in the development and metamorphosis of insects [96]. In the fruit fly *D. melanogaster* and the moths *B. mori* and *M. sexta*, the 26-hydroxylase encoded by *CYP18A1* affects developmental processes by inactivating 20E and degrading it to 20,26-dihydroxyecdysone through the addition of an OH group to the 26th carbon of the hormone molecule [96,97]. Our study revealed homologues of *CYP18A1* in all major panarthropod subgroups, including the onychophoran *E. rowelli*, in which CYP18A1 might play a similar role in 20E degradation as this ecdysteroid has been demonstrated in the closely related species *E. leuckartii* [25]. By contrast, we found no unambiguous homologues of *CYP18A1* in available genomes from nematodes and the priapulid *P. caudatus*, suggesting that this gene might have been recruited for degradation of 20E (or a related hormone) in the last common ancestor of Panarthropoda.

## 5. Conclusion

To clarify whether the early moulting pathway of the fruit fly *D. melanogaster* is a conserved feature of Ecdysozoa, we analysed candidate genes of this pathway in various bilaterians. While the homologues of genes encoding the heterodimer receptor complex EcR/USP and the Early genes are present in all major ecdysozoan subgroups, to our surprise we found only a few genes of the canonical ecdysteroid pathway in these taxa. Although our results revealed no homologues of the Rieske-domain gene *nvd* in onychophorans, tardigrades and priapulids, this gene might have been present in the last common ancestor of Ecdysozoa and was most likely lost in these three ecdysozoan subgroups (figure 6). Hence, it is unclear whether the dehydrogenation of cholesterol to 7-dehydrocholesterol occurs in these animals and, if so, which enzyme accomplishes this dehydrogenation. Similarly, we found homologues of *CYP18A1*—a gene belonging to the degradation pathway of 20E—only in panarthropods; it is, therefore, unclear which gene was responsible for the degradation of ecdysteroids in the last common ancestor of Ecdysozoa.

Another unexpected result of our study is the finding that most likely none of the Halloween genes were part of the ecdysteroid pathway in the last common ancestor of Ecdysozoa, and that only the Halloween

gene *sad* was present in the last common ancestor of Panarthropoda (figure 6). Although functional assays would be required to clarify whether or not this gene is involved in the hydroxylation of ecdysone to 2-deoxyecdysone in onychophorans and tardigrades, our results clearly indicate that *sad* was the first Halloween gene to evolve in the panarthropod lineage, whereas the remaining Halloween genes might have originated subsequently in arthropods by gene duplications within the cytochrome P450 superfamily.

Taken together, these results suggest that most of the key players responsible for the conversion of cholesterol to 20E (or a related hormone) in the last common ancestor of Ecdysozoa are currently unknown. This is astonishing, given that ecdysis—a process which is believed to be governed by the ecdysteroid hormones—is regarded as the most prominent autapomorphy of this clade [3–5]. Future studies should, therefore, focus on aspects of the endocrine control of this process in understudied ecdysozoan subgroups, to gain a full understanding of this vital process in this important clade.

**Ethics.** Permission for the *Euperipatoides rowelli* (Onychophora) collection was obtained from the National Parks & Wildlife Service New South Wales (permit no. SL100159). Permission for the export of specimens was obtained from the Department of Sustainability, Environment, Water, Population and Communities (permit no. PWSP104061). Specimens of the eutardigrade species *Hypsibius exemplaris* (Tardigrada) were obtained commercially from Sciento (Manchester, UK).

**Data accessibility.** All data are available in the electronic supplementary material.

**Authors' contributions.** G.M. and I.S. designed the experiments. I.S. and N.K. carried out genome and transcriptome analyses. I.S., N.K. and L.H. performed phylogenetic analyses. All authors discussed the results and wrote the manuscript.

**Competing interests.** The authors declare that they have no conflict of interest.

**Funding.** This research was supported by the Hong Kong Research Grant Council ECS grant (24100015), CUHK Direct grant (no. 4053096) and CUHK Faculty of Science Strategic Development Scheme to J.H., and the German Research Foundation (DFG: Ma 4147/3-1) to G.M.

**Acknowledgements.** Pre-publication access to genomic data from *E. rowelli* generated as part of the i5 K pilot at the Baylor College of Medicine Human Genome Sequencing Center was kindly provided by Stephen Richards. We further acknowledge Richard K. Wilson, Director, and the McDonnell Genome Institute, Washington School of Medicine for providing the *Priapulius caudatus* 5.0 sequence assembly (AXZU00000000.2). We are thankful to the members of the Mayer laboratory for their support with animal husbandry and to Martin Schlegel and members of his group for general support. We gratefully acknowledge Paul M. Whittington for his assistance with collecting the specimens, and Noel N. Tait for his help with permits. The staffs of the National Parks & Wildlife Service of New South Wales and the Department of Sustainability, Environment, Water, Population and Communities are gratefully acknowledged for providing the collection and export permits. Vladimir Gross, Stefan Schaffer and Justus Hennecke provided useful suggestions on the manuscript.

## References

- Giribet G, Edgecombe GD. 2017 Current understanding of Ecdysozoa and its internal phylogenetic relationships. *Integr. Comp. Biol.* **57**, 455–466. (doi:10.1093/icb/ixc072)
- Kocot KM. 2016 On 20 years of Lophotrochozoa. *Org. Divers. Evol.* **16**, 329–343. (doi:10.1007/s13127-015-0261-3)
- Aguinaldo AMA, Turbeville JM, Linford LS, Rivera MC, Garey JR, Raff RA, Lake JA. 1997 Evidence for a clade of nematodes, arthropods and other moulting animals. *Nature* **387**, 489–493. (doi:10.1038/387489a0)
- Schmidt-Rhaesa T, Lemburg C, Ehlers U, Garey JR, Bartolomaeus A. 1998 The position of the Arthropoda in the phylogenetic system. *J. Morphol.* **238**, 263–285. (doi:10.1002/(SICI)1097-4687(199812)238:3<<263::AID-JMOR1>>3.0.CO;2-L)
- Nielsen C. 2012 Ecdysozoa. In *Animal evolution: interrelationships of the living phyla* (ed. C Nielsen), 3rd edn, p. 249. Oxford, UK: Oxford University Press.
- Rota-Stabelli O, Daley AC, Pisani D. 2013 Molecular timetrees reveal a Cambrian colonization of land and a new scenario for ecdysozoan evolution. *Curr. Biol.* **23**, 392–398. (doi:10.1016/j.cub.2013.01.026)
- Ewer J. 2005 How the ecdysozoan changed its coat. *PLoS Biol.* **3**, e349. (doi:10.1371/journal.pbio.0030349)
- Nijhout HF. 2013 Arthropod developmental endocrinology. In *Arthropod biology and evolution* (ed. G Minelli, G Fusco), pp. 123–148. Heidelberg, Germany: Springer.
- Krishnakumar A, Schneiderman HA. 1970 Control of molting in mandibulate and chelicerate arthropods by ecdysones. *Biol. Bull.* **139**, 520–538. (doi:10.2307/1540371)
- Karlson P. 1996 On the hormonal control of insect metamorphosis: a historical review. *Int. J. Dev. Biol.* **40**, 93–96.
- Hampshire F, Horn DHS. 1966 Structure of crustecdysone, a crustacean moulting hormone. *Chem. Comm. (London)* **2**, 37–38. (doi:10.1039/C19660000037)
- Rota-Stabelli O, Kayal E, Gleeson D, Daub J, Boore JL, Telford MJ, Pisani D, Blaxter M, Lavrov DV. 2010 Ecdysozoan mitogenomics: evidence for a common origin of the legged invertebrates, the Panarthropoda. *Genome Biol. Evol.* **2**, 425–440. (doi:10.1093/gbe/evq030)
- Edgar BA. 2006 How flies get their size: genetics meets physiology. *Nat. Rev. Genet.* **7**, 907–916. (doi:10.1038/nrg1989)
- Igarashi F, Ogihara MH, Iga M, Kataoka H. 2018 Cholesterol internalization and metabolism in insect prothoracic gland, a steroidogenic organ, via lipoproteins. *Steroids* **134**, 110–116. (doi:10.1016/j.steroids.2018.01.012)
- Warren JT, Yerushalmi Y, Shimell MJ, O'Connor MB, Restifo LL, Gilbert LI. 2006 Discrete pulses of molting hormone, 20-hydroxyecdysone, during late larval development of *Drosophila melanogaster*: correlations with changes in gene activity. *Dev. Dyn.* **235**, 315–326. (doi:10.1002/dvdy.20626)
- Yoshiyama T, Namiki T, Mita K, Kataoka H, Niwa R. 2006 Neverland is an evolutionarily conserved Rieske-domain protein that is essential for ecdysone synthesis and insect growth. *Development* **133**, 2565–2574. (doi:10.1242/dev.02428)

17. Yoshiyama-Yanagawa T *et al.* 2011 The conserved Rieske oxygenase DAF-36/Neverland is a novel cholesterol-metabolizing enzyme. *J. Biol. Chem.* **286**, 25 756–25 762. (doi:10.1074/jbc.M111.244384)
18. Ono H *et al.* 2006 *Spook* and *Spookier* code for stage-specific components of the ecdysone biosynthetic pathway in Diptera. *Dev. Biol.* **298**, 555–570. (doi:10.1016/j.ydbio.2006.07.023)
19. Rewitz KF, O'Connor MB, Gilbert LI. 2007 Molecular evolution of the insect Halloween family of cytochrome P450s: phylogeny, gene organization and functional conservation. *Insect Biochem. Mol. Biol.* **37**, 741–753. (doi:10.1016/j.ibmb.2007.02.012)
20. Niwa R, Niwa YS. 2014 Enzymes for ecdysteroid biosynthesis: their biological functions in insects and beyond. *Biosci. Biotechnol. Biochem.* **78**, 1283–1292. (doi:10.1080/09168451.2014.942250)
21. Zhu XX, Oliver JH, Dotson EM. 1991 Epidermis as the source of ecdysone in an argasid tick. *Proc. Natl Acad. Sci. USA* **88**, 3744–3747. (doi:10.1073/pnas.88.9.3744)
22. Petryk A *et al.* 2003 Shade is the *Drosophila* P450 enzyme that mediates the hydroxylation of ecdysone to the steroid insect molting hormone 20-hydroxyecdysone. *Proc. Natl Acad. Sci. USA* **100**, 13 773–13 778. (doi:10.1073/pnas.2336088100)
23. Yao TP, Forman BM, Jiang Z, Cherbas L, Chen JP, McKeown M, Cherbas P, Evans RM. 1993 Functional ecdysone receptor is the product of *Ecr* and *ultraspiracle* genes. *Nature* **366**, 476. (doi:10.1038/366476a0)
24. Billas IML, Moulinier L, Rochel N, Moras D. 2001 Crystal structure of the ligand-binding domain of the Ultraspiracle protein UsP, the ortholog of retinoid X receptors in insects. *J. Biol. Chem.* **276**, 7465–7474. (doi:10.1074/jbc.M008926200)
25. Puthumana J, Lee MC, Han J, Kim HS, Hwang DS, Lee JS. 2017 *Ecdysone receptor* (Ecr) and *ultraspiracle* (USP) response to water accommodated fractions (WAFs). *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* **192**, 7–15. (doi:10.1016/j.cbpc.2016.11.002)
26. Ashburner M, Richards G. 1976 Sequential gene activation by ecdysone in polytene chromosomes of *Drosophila melanogaster*: III. Consequences of ecdysone withdrawal. *Dev. Biol.* **54**, 241–255. (doi:10.1016/0012-1606(76)90302-X)
27. Segraves WA, Hogness DS. 1990 The *E75* ecdysone-inducible gene responsible for the 75B early puff in *Drosophila* encodes two new members of the steroid receptor superfamily. *Genes Dev.* **4**, 204–219. (doi:10.1101/gad.4.2.204)
28. Li TR, White KP. 2003 Tissue-specific gene expression and ecdysone-regulated genomic networks in *Drosophila*. *Dev. Cell* **5**, 59–72. (doi:10.1016/S1534-5807(03)00192-8)
29. King-Jones K, Thummel CS. 2005 Nuclear receptors—a perspective from *Drosophila*. *Nat. Rev. Genet.* **6**, 311. (doi:10.1038/nrg1581)
30. Baldwin WS, Marko PB, Nelson DR. 2009 The cytochrome P450 (CYP) gene superfamily in *Daphnia pulex*. *BMC Genomics* **10**, 169. (doi:10.1186/1471-2164-10-169)
31. Cheong SPS, Huang J, Bendena WG, Torbe SS, Hui JHL. 2015 Evolution of ecdysis and metamorphosis in arthropods: the rise of regulation of juvenile hormone. *Integr. Comp. Biol.* **55**, 878–890. (doi:10.1093/ich/icc066)
32. Cao JQ, Tong WS, Yu HY, Tobe SS, Bendena WG, Hui JHL. 2017 The role of microRNAs in *Drosophila* regulation of insulin-like peptides and ecdysteroid signalling: where are we now? In *Insect Epigenetics* (ed. HBT Verlinden), pp. 55–85. New York, NY: Academic Press.
33. Martin C, Gross V, Hering L, Tepper B, Jahn H, de Sena Oliveira I, Stevenson PA, Mayer G. 2017 The nervous and visual systems of onychophorans and tardigrades: learning about arthropod evolution from their closest relatives. *J. Comp. Physiol. A* **203**, 565–590. (doi:10.1007/s00359-017-1186-4)
34. Hoffmann KH. 1997 Ecdysteroids in adult females of a 'walking worm': *Euperipatoides leuckartii* (Onychophora, Peripatopsidae). *Invertebr. Reprod. Dev.* **32**, 27–30. (doi:10.1080/07924259.1997.9672601)
35. Cleator M, Delves CJ, Howells RE, Rees HH. 1987 Identity and tissue localization of free and conjugated ecdysteroids in adults of *Dirofilaria immitis* and *Ascaris suum*. *Mol. Biochem. Parasitol.* **25**, 93–105. (doi:10.1016/0166-6851(87)90022-3)
36. Shea C, Hough D, Xiao J, Tertzinis G, Maina CV. 2004 An *rxr/usp* homolog from the parasitic nematode, *Dirofilaria immitis*. *Gene* **324**, 171–182. doi:10.1016/j.gene.2003.09.032
37. Frand AR, Russel S, Ruvkun G. 2005 Functional genomic analysis of *C. elegans* molting. *PLoS Biol.* **3**, e312. (doi:10.1371/journal.pbio.0030312)
38. Lažetić V, Fay DS. 2017 Molting in *C. elegans*. *Worm* **6**, e1330246. (doi:10.1080/21624054.2017.1330246)
39. Markov GV, Tavares R, Dauphin-Villemant C, Demeneix BA, Baker ME, Laudet V. 2009 Independent elaboration of steroid hormone signaling pathways in metazoans. *Proc. Natl Acad. Sci. USA* **106**, 11 913–11 918. (doi:10.1073/pnas.0812138106)
40. Mendis AHW, Rees HH, Goodwin TW. 1984 The occurrence of ecdysteroids in the cestode, *Moniezia expansa*. *Mol. Biochem. Parasitol.* **10**, 123–138. (doi:10.1016/0166-6851(84)90001-X)
41. Nolte A, Koolman J, Dorlöchter M, Straub H. 1986 Ecdysteroids in the dorsal bodies of pulmonates (Gastropoda): synthesis and release of ecdysone. *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* **84**, 777–782. (doi:10.1016/0300-9629(86)90405-6)
42. García M, Gharbi J, Girault JP, Hetru C, Lafont R. 1989 Ecdysteroid metabolism in leeches. *Invertebr. Reprod. Dev.* **15**, 57–68. (doi:10.1080/07924259.1989.9672022)
43. Paxton H. 2005 Molting polychaete jaws—ecdysozoans are not the only molting animals. *Evol. Dev.* **7**, 337–340. (doi:10.1111/j.1525-142X.2005.05039.x)
44. Pilato G *et al.* 2005 The clade Ecdysozoa, perplexities and questions. *Zool. Anz.* **244**, 43–50. (doi:10.1016/j.jcz.2005.04.001)
45. Sauber F, Reuland M, Berchtold JP, Hetru C, Tsoupras G, Luu B, Moritz ME, Hoffmann JA. 1983 Molting cycle and ecdysteroids in the leech, *Hirudo medicinalis*. *C. R. Acad. Sci. - Series III* **296**, 413.
46. Barker GC, Chitwood DJ, Rees HH. 1990 Ecdysteroids in helminths and annelids. *Invertebr. Reprod. Dev.* **18**, 1–11. (doi:10.1080/07924259.1990.9672124)
47. The *C. elegans* Sequencing Consortium. 1998 Genome sequence of the nematode *C. elegans*: a platform for investigating biology. *Science* **282**, 2012–2018. (doi:10.1126/science.282.5396.2012)
48. Koutsovoulos G *et al.* 2015 The genome of the tardigrade *Hypsibius dujardini*. *bioRxiv*. (doi:10.1101/033464)
49. Hering L, Mayer G. 2014 Analysis of the opsin repertoire in the tardigrade *Hypsibius dujardini* provides insights into the evolution of opsin genes in Panarthropoda. *Genome Biol. Evol.* **6**, 2380–2391. (doi:10.1093/gbe/evu193)
50. Hashimoto T *et al.* 2016 Extremotolerant tardigrade genome and improved radiotolerance of human cultured cells by tardigrade-unique protein. *Nat. Commun.* **7**, 12808. (doi:10.1038/ncomms12808)
51. Hering L *et al.* 2012 Opsins in Onychophora (velvet worms) suggest a single origin and subsequent diversification of visual pigments in arthropods. *Mol. Biol. Evol.* **29**, 3451–3458. (doi:10.1093/molbev/mss148)
52. Hill CA, Wikel SK 2005 The *Ixodes scapularis* Genome Project: an opportunity for advancing tick research. *Trends Parasitol.* **21**, 151–153. (doi:10.1016/j.pt.2005.02.004)
53. Grbić M *et al.* 2011 The genome of *Tetranychus urticae* reveals herbivorous pest adaptations. *Nature* **479**, 487. (doi:10.1038/nature10640)
54. Sanggaard KW *et al.* 2014 Spider genomes provide insight into composition and evolution of venom and silk. *Nat. Commun.* **5**, 3765. (doi:10.1038/ncomms4765)
55. Chipman AD *et al.* 2014 The first myriapod genome sequence reveals conservative arthropod gene content and genome organisation in the centipede *Strigamia maritima*. *PLoS Biol.* **12**, e1002005. (doi:10.1371/journal.pbio.1002005)
56. Baer A, Mayer G. 2012 Comparative anatomy of slime glands in Onychophora (velvet worms). *J. Morphol.* **273**, 1079–1088. (doi:10.1002/jmor.20044)
57. Beckmann H, Hering L, Henze MJ, Kelber A, Stevenson PA, Mayer G. 2015 Spectral sensitivity in Onychophora (velvet worms) revealed by electroretinograms, phototactic behaviour and opsin gene expression. *J. Exp. Biol.* **218**, 915–922. (doi:10.1242/jeb.116780)
58. Gąsiorek P, Stec D, Morek W, Michalczyk Ł. 2018 An integrative redescription of *Hypsibius dujardini* (Doyère, 1840), the nominal taxon for Hypsibioidae (Tardigrada: Eutardigrada). *Zootaxa* **4415**, 45–75. (doi:10.11646/zootaxa.4415.1.2)
59. Gross V, Mayer G. 2015 Neural development in the tardigrade *Hypsibius dujardini* based on anti-acetylated  $\alpha$ -tubulin immunolabeling.

- Evol. Dev.* **6**, 12. (doi:10.1186/s13227-015-0008-4)
60. Gross V, Minich I, Mayer G. 2017 External morphogenesis of the tardigrade *Hypsibius dujardini* as revealed by scanning electron microscopy. *J. Morphol.* **278**, 563–573. (doi:10.1002/jmor.20654)
61. Gross V, Bährle R, Mayer G. 2018 Detection of cell proliferation in adults of the water bear *Hypsibius dujardini* (Tardigrada) via incorporation of a thymidine analog. *Tissue Cell* **51**, 77–83. (doi.org/10.1016/j.tice.2018.03.005)
62. Boothby TC *et al.* 2015 Evidence for extensive horizontal gene transfer from the draft genome of a tardigrade. *Proc. Natl Acad. Sci. USA* **112**, 15 976–15 981. (doi:10.1073/pnas.1510461112)
63. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997 Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**, 3389–3402. (doi:10.1093/nar/25.17.3389)
64. Kersey PJ *et al.* 2016 Ensembl Genomes 2016: more genomes, more complexity. *Nucleic Acids Res.* **44**, D574–D580. (doi:10.1093/nar/gkv1209)
65. McWilliam H, Li W, Uludag M, Squizzato S, Park YM, Buso N, Cowley AP, Lopez R. 2013 Analysis tool web services from the EMBL-EBI. *Nucleic Acids Res.* **41**, W597–W600. (doi:10.1093/nar/gkt376)
66. Finn RD *et al.* 2016 The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Res.* **44**, D279–D285. (doi:10.1093/nar/gkv1344)
67. Schultz J, Milpetz F, Bork P, Ponting CP. 1998 SMART, a simple modular architecture research tool: identification of signaling domains. *Proc. Natl Acad. Sci. USA* **95**, 5857–5864. (doi:10.1073/pnas.95.11.5857)
68. Sharrocks AD, Brown AL, Ling Y, Yates PR. 1997 The ETS-domain transcription factor family. *Int. J. Biochem. Cell Biol.* **29**, 1371–1387. (doi:10.1016/S1357-2725(97)00086-1)
69. Katoh K, Standley DM. 2013 MAFFT Multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780. (doi:10.1093/molbev/mst010)
70. Stamatakis A. 2006 RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* **22**, 2688–2690. (doi:10.1093/bioinformatics/btl446)
71. Letunic I, Bork P. 2016 Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Res.* **44**, W242–W245. (doi:10.1093/nar/gkw290)
72. Lafont R. 1991 Reverse endocrinology, or ‘hormones’ seeking functions. *Insect Biochem.* **21**, 697–721. (doi:10.1016/0020-1790(91)90112-R)
73. Grieneisen ML. 1994 Recent advances in our knowledge of ecdysteroid biosynthesis in insects and crustaceans. *Insect Biochem. Mol. Biol.* **24**, 115–132. (doi:10.1016/0965-1748(94)90078-7)
74. Chavez VM, Marques G, Delbecq JP, Kobayashi K, Hollingsworth M, Burr J, Natzle JE, O’Connor MB. 2000 The *Drosophila disembodied* gene controls late embryonic morphogenesis and codes for a cytochrome P450 enzyme that regulates embryonic ecdysone levels. *Development* **127**, 4115–4126.
75. Rewitz KF, Rycyzynski R, Warren JT, Gilbert LI. 2006 The Halloween genes code for cytochrome P450 enzymes mediating synthesis of the insect moulting hormone. *Biochem. Soc. Trans.* **34**, 1256–1260. (doi:10.1042/bst0341256)
76. Nüsslein-Volhard C, Wieschaus E, Kluding H. 1984 Mutations affecting the pattern of the larval cuticle in *Drosophila melanogaster*. *Roux Arch. Dev. Biol.* **193**, 267–282. (doi:10.1007/BF00848156)
77. Cabrera AR, Shirk PD, Evans JD, Hung K, Sims J, Alborn H, Teal PEA. 2015 Three Halloween genes from the Varroa mite, *Varroa destructor* (Anderson & Trueman) and their expression during reproduction. *Insect Mol. Biol.* **24**, 277–292. (doi:10.1111/imb.12155)
78. Aguilaniu H, Fabrizio P, Witting M. 2016 The role of daftachronic acid signaling in development and longevity in *Caenorhabditis elegans*: digging deeper using cutting-edge analytical chemistry. *Front. Endocrinol.* **7**, 12. (doi:10.3389/fendo.2016.00012)
79. Fleming MW. 1985 *Ascaris suum*: role of ecdysteroids in molting. *Exp. Parasitol.* **60**, 207–210. (doi:10.1016/0014-4894(85)90024-4)
80. Parihar M, Minton RL, Flowers S, Holloway A, Morehead BE, Paille J, Gissendammer CR. 2010 The genome of the nematode *Pristionchus pacificus* encodes putative homologs of RXR/Usp and Ecr. *Gen. Comp. Endocrinol.* **167**, 11–17. (doi:10.1016/j.ygcen.2010.02.005)
81. Tzertzinis G, Egaña AL, Palli SR, Robinson-Rechavi M, Gissendanner CR, Liu C, Unnash TR, Maina CV. 2010 Molecular evidence for a functional ecdysone signaling system in *Brugia malayi*. *PLoS Negl. Trop. Dis.* **4**, e625. (doi:10.1371/journal.pntd.0000625)
82. Qu Z, Kenny NJ, Lam HM, Chan TF, Chu KH, Bendena WG, Tobe SS, Hui JHL. 2015 How did arthropod sesquiterpenoids and ecdysteroids arise? Comparison of hormonal pathway genes in non-insect arthropod genomes. *Genome Biol. Evol.* **7**, 1951–1959. (doi:10.1093/gbe/evv120)
83. Oro AE, McKeown M, Evans RM. 1992 The *Drosophila* retinoid X receptor homolog *ultraspiracle* functions in both female reproduction and eye morphogenesis. *Development* **115**, 449–462.
84. Martín D, Wang S-F, Raikhel AS. 2001 The vitellogenin gene of the mosquito *Aedes aegypti* is a direct target of ecdysteroid receptor. *Mol. Cell. Endocrinol.* **173**, 75–86. (doi:10.1016/S0303-7207(00)00413-5)
85. Xu J, Tan A, Palli SR. 2010 The function of nuclear receptors in regulation of female reproduction and embryogenesis in the red flour beetle, *Tribolium castaneum*. *J. Insect Physiol.* **56**, 1471–1480. (doi:10.1016/j.jinsphys.2010.04.004)
86. Talbot WS, Swyryd EA, Hogness DS. 1993 *Drosophila* tissues with different metamorphic responses to ecdysone express different ecdysone receptor isoforms. *Cell* **73**, 1323–1337. (doi:10.1016/0092-8674(93)90359-X)
87. Jindra M, Malone F, Hiruma K, Riddiford LM. 1996 Developmental profiles and ecdysteroid regulation of the mRNAs for two ecdysone receptor isoforms in the epidermis and wings of the tobacco hornworm, *Manduca sexta*. *Dev. Biol.* **180**, 258–272. (doi:10.1006/dbio.1996.0299)
88. Riddiford LM, Hiruma K, Zhou X, Nelson CA. 2003 Insights into the molecular basis of the hormonal control of molting and metamorphosis from *Manduca sexta* and *Drosophila melanogaster*. *Insect Biochem. Mol. Biol.* **33**, 1327–1338. (doi:10.1016/j.ibmb.2003.06.001)
89. Dubrovsky EB. 2005 Hormonal cross talk in insect development. *Trends Endocrinol. Metab.* **16**, 6–11. (doi:10.1016/j.tem.2004.11.003)
90. Zhao X, Qin Z, Liu W, Liu X, Moussian B, Ma E, Li S, Zhang J. 2017 Nuclear receptor HR3 controls locust molt by regulating chitin synthesis and degradation genes of *Locusta migratoria*. *Insect Biochem. Mol. Biol.* **92**, 1–11. (doi:10.1016/j.ibmb.2017.11.001)
91. Stone BL, Thummel CS. 1993 The *Drosophila* 78C early late puff contains *E78*, an ecdysone-inducible gene that encodes a novel member of the nuclear hormone receptor superfamily. *Cell* **75**, 307–320. (doi:10.1016/0092-8674(93)80072-m)
92. Yao Q, Zhang D, Tang B, Chen J, Chen J, Lu L, Zhang W. 2010 Identification of 20-hydroxyecdysone late-response genes in the chitin biosynthesis pathway. *PLoS ONE* **5**, e14058. (doi:10.1371/journal.pone.0014058)
93. Song Y, Villeneuve DL, Toyota K, Iguchi T, Tollefsen KE. 2017 Ecdysone receptor agonism leading to lethal molting disruption in arthropods: review and adverse outcome pathway development. *Environ. Sci. Technol.* **51**, 4142–4157. (doi:10.1021/acs.est.7b00480)
94. Kostrouchova M, Krause M, Kostrouch Z, Rall JE. 2001 Nuclear hormone receptor CHR3 is a critical regulator of all four larval molts of the nematode *Caenorhabditis elegans*. *Proc. Natl Acad. Sci. USA* **98**, 7360–7365. (doi:10.1073/pnas.131171898)
95. Gissendanner CR, Crossgrove K, Kraus KA, Maina CV, Sluder AE. 2004 Expression and function of conserved nuclear receptor genes in *Caenorhabditis elegans*. *Dev. Biol.* **266**, 399–416. (doi:10.1016/j.ydbio.2003.10.014)
96. Guittard E, Blais C, Maria A, Parvy JP, Pasricha S, Lumb C, Lafont R, Daborn PJ, Dauphin-Villemant C. 2011 CYP18A1, a key enzyme of *Drosophila* steroid hormone inactivation, is essential for metamorphosis. *Dev. Biol.* **349**, 35–45. (doi:10.1016/j.ydbio.2010.09.023)
97. Rewitz KF, Gilbert LI. 2008 *Daphnia* Halloween genes that encode cytochrome P450s mediating the synthesis of the arthropod molting hormone: evolutionary implications. *BMC Evol. Biol.* **8**, 60. (doi:10.1186/1471-2148-8-60)