

Boxwood Borer *Heterobostrychus brunneus* (Coleoptera: Bostrichidae) Infesting Dried Cassava: A Current Record from Southern Ethiopia

Aditya Parmar,^{1,2} Sascha M. Kirchner,¹ Henning Langguth,³ Thomas F. Döring,⁴ and Oliver Hensel¹

¹Department of Agricultural and Biosystems Engineering, University of Kassel, Nordbahnhofstrasse 1a, Witzenhausen 37213, Germany,

²Corresponding author, e-mail: aditya.parmar@daad-alumni.de, ³Institute for Biology, University of Kassel, Heinrich-Plett-Strasse 40, Kassel 34132, Germany, and ⁴Department of Agronomy and Crop Science, Humboldt University Berlin, Albrecht-Thaer Weg 5, Berlin 14195, Germany

Subject Editor: Nickolas Kavallieratos

Received 30 June 2016; editorial decision 1 November 2016

Abstract

Insect specimens of adult beetles and larvae of 7–9 and 9–10 mm length, respectively were collected from infested dry cassava at two locations from multiple stores in southern Ethiopia. The specimens were identified as *Heterobostrychus brunneus* (Murray, 1867) commonly known as boxwood borer and auger beetle. The study presents a current record of *H. brunneus* in Ethiopia, particularly in the context of infesting food products. Additionally, a wide geographical distribution of the pest was reviewed and presented in this article. Current evidence suggests that *H. brunneus* is a serious pest of forest wood, structural timbers, and dried food products and that it carries a risk to be introduced into various other parts of the world via global trade.

Key words: Cassava, COI, Ethiopia, *Heterobostrychus brunneus*

The genus *Heterobostrychus* Lesne, 1899 belongs to the Bostrichidae family and comprises species that are commonly considered as forest wood pests that can also infest foodstuff and household structural timbers. Most of the bostrichids are known to be polyphagous in nature; both adult and larvae feed on a variety of woods and obtain their nutrition from starch and sugars present in their hosts. Moreover, some species can also feed on food products (Hill 2002, Liu et al. 2008). Species belonging to *Heterobostrychus* are notorious pests of significant economic importance in tropical and subtropical regions and are categorized as powder-post beetle because of their peculiar nature of reducing the wood to powdery dust (Findlay 1985, Liu et al. 2008, Azmi et al. 2011). Six known species of the genus are *H. aequalis* (Waterhouse, 1884), *H. ambigenus* (Lesne, 1920), *H. brunneus* (Murray, 1867), *H. hamatipennis* (Lesne, 1895), *H. pileatus* (Lesne, 1899), and *H. unicornis* (Waterhouse, 1879) (Beiriger 2010, ITIS 2016). Some species of the genus (*H. aequalis*, *H. brunneus* and *H. hamatipennis*) have regularly been introduced in various parts of the world (Borowski and Wegrzynowicz 2007) due to increased global trade in timber and wood products. According to Azmi et al. (2011), one of the species (*H. aequalis*) was introduced most likely via trade to various countries, including Belgium, Canada, France, Germany, Israel, Italy, Japan, New Zealand, and the United States.

H. brunneus is commonly called boxwood borer due to its presence in hardwood packings. This habit makes it a quarantine pest in many countries where the species is non-endemic or not yet established (Wylie et al. 2008). Adults and larvae of *H. brunneus* can affect timber in green condition and after seasoning (Findlay 1985). The beetle is about 6–13 mm long and dark reddish brown to black in color. Full grown larvae are pale yellow and about 7–10 mm long (Fisher 1950, Robinson 2005). Booth et al. (1990) describe *H. brunneus* as being similar to *H. aequalis* but having recumbent hair and an absence of tubercles on the top margins of the elytral declivity.

Cassava (*Manihot esculenta* Crantz) is an important root crop and is cultivated in various parts of Africa, Asia, and South America. Globally, cassava is the sixth largest crop after wheat, rice, maize, potato, and barley (Lebot 2009). Estimates suggest it is a staple food for more than 800 million people, mostly living in the least industrialized tropical and sub-tropical regions of the world (Howeler et al. 2013). However, the subsistence status of this crop species is rapidly changing to a cash crop with its enhanced usage in multiple industrial applications such as the production of paper, textile, plywood, glue, biofuel, animal feed and beverage. Cassava in Ethiopia is relatively new, it was introduced in 1960, but its importance dramatically increased in the country after the famine of 1984 (Kebede et al. 2012). Presently, it is an important crop in the south




Fig. 1. Sites from where specimens of the adult and larvae (*H. brunneus*) were collected [SNNP (Southern Nations, Nationalities, and Peoples' Region). Bale - N6° 55.175' E37° 32.075', Elevation: 1245m asl; Sodo - N6° 51.481' E37° 46.141'. Elevation: 2017m asl.

and south-western parts of Ethiopia where the local population tends to rely more on root crops.

During storage, dried cassava is highly susceptible to a variety of storage insects and pests. The damage is intensified by tropical conditions resulting in significant weight loss in a short time (Hodges et al. 1985). Stumpf (1998) reported *H. brunneus* (Ghana) as a destructive pest of dried cassava. The species not only damages the dried products, but it has also been reported as a common stem borer for standing cassava plantations (Bellotti and van Schoonhoven 1978). Hence, the presence of *H. brunneus* poses a dual threat to cassava during production and storage and requires immediate attention for control and prevention. The purpose of this study is to provide details on the presence of *H. brunneus* infestation in the dried cassava chips and chunks during storage in southern Ethiopia. Key morphological features and DNA barcode of the species are reported. Additionally, a global distribution data for the species based on current information was generated.

Material and Methods

Location and Survey

A field survey to assess the post-harvest losses and related causes in the cassava value chain was conducted in Wolayita Zone of Southern Region (SNNPR) Ethiopia in October–December 2015. In the course of this study, 60 traders were surveyed by administering semi-structured questionnaires and conducting store inspection using pictographic illustrations of the main storage pests in three

districts. Along with other more common insect species, unidentified black, brownish beetles (adults of 8–9 mm length) were presented and reported by traders infesting dried cassava chips at their storage systems. Sodo and Bale town in Sodo Zuriya and Kindo Koyisha districts, respectively were the two locations in the southern highland of Ethiopia from where the specimens were collected. The climate of Sodo is classified as warm and temperate (Köppen-Geiger: *Cfb*) with a mean annual temperature of 19.3°C and 1,484 mm of annual precipitation (Gonfa 1996, Calimate Data Org 2016). Bale has a tropical climate (Köppen-Geiger: *Aw*) with a mean annual temperature and an annual precipitation of 22.7°C and 1,200 mm, respectively (Gonfa 1996, Calimate Data Org 2016). The exact coordinates and elevations of the locations are presented in Figure 1.

The specimens of the unidentified adult beetles along with larvae were collected from the storage houses and kept in 70% ethanol until identification. The species was identified using several keys for adults (Fisher 1950, Walker 2005, Beiriger 2010) and larvae (Lesne 1924, Robinson 2005, Schabel 2006). The online source the Integrated Taxonomic Information System (ITIS 2016) and the Department of Entomology, Iowa State University the BugGuide (2016) were used for nomenclature classification.

Photographs of the specimens (Figs. 3–5) were taken with a stereomicroscope Leica-EZ4HD (Leica Microsystems, Wetzlar, Germany). DNA barcoding of the adult beetle was conducted, which was based on a 658-bp long region of the mitochondrial gene for cytochrome c oxidase I (COI) amplified using primers COL6 (5'- TYTCHACA

Table 1. Distribution of *H. brunneus* (Murray, 1867) based on current information.

Region	Status	Host	References
Africa			
Burundi	Present	NI*	Hagstrum and Subramanyam (2009)
Cameroon	Present	NI	Vrydagh (1960), Alene et al. (2005), EPPO (2016a)
Cape Verde	Present	Timber	Fisher (1950), EPPO (2016b)
Congo	Present	Logs	Fisher (1950), Borowski and Wegrzynowicz (2007), GBIF (2016)
Egypt	Present	NI	Borowski (2007), Hagstrum and Subramanyam (2009)
Ethiopia (Abyssinia)	Present	NI	Vrydagh (1960)
Gambia	Present	NI	GBIF (2016)
Ghana	Present	Cassava, sapwood, lumber	Ashiru (1989), Stumpf (1998), Wagner et al. (2008)
Ivory coast	Present	Wood	Becker (1980)
Kenya	Present	NI	Wasonga et al. (2015)
Madagascar	Present	NI	Fisher (1950)
Mozambique	Present	NI	Vrydagh (1955b)
Nigeria	Present	Lumber, dead logs	Roberts (1967), Ashiru (1989), Borowski and Wegrzynowicz (2007)
Rawanda	Present	NI	Hagstrum and Subramanyam (2009)
Senegal	Present	NI	GBIF (2016)
Seychelles Islands	Present	Timber	Fisher (1950), EPPO (2016b)
South Africa	Present	Bamboo, timber	Lesne (1924), Pringle (1938), Fisher (1950), GBIF (2016)
Sudan	Present	NI	Hagstrum and Subramanyam (2009)
The Republic of Guinea	Present	Potato tubers	Fisher (1950)
Tanzania	Present	Forest wood	Madoffe et al. (2000), Schabel (2006), GBIF (2016)
Togo	Present	NI	Vrydagh (1955a)
Uganda	Present	Dried sweet potato,	Davies (1960), Mwanga and Wanyera (1988)
Zambia	Present	Sawn Timber, Logs	Selander and Pearce (1984), Lemmens (2012)
Zimbabwe	Present	NI	Walker (2005), GBIF (2016)
Europe			
Belgium	Intercepted	NI	Borowski (2007), Hagstrum and Subramanyam (2009)
United Kingdom	Intercepted	NI	Borowski (2007), Hagstrum and Subramanyam (2009), Fauna Europaea (2016)
France	Intercepted	NI	Borowski (2007), Hagstrum and Subramanyam (2009), Brustel and Aberlenc (2014)
Germany	Intercepted	Imported timber or bamboo	Cymorek 1970, Borowski (2007), Fauna Europaea (2016), GBIF (2016)
Italy	Intercepted	Wood and wood products	Gambetta and Orlandi (1982), Audisio et al. (1995), Ratti (2004), Borowski (2007), Fauna Europaea (2016), GBIF (2016)
Poland	Intercepted	NI	Hagstrum and Subramanyam (2009), Borowski (2007)
Portugal	Intercepted	NI	Baena and Zuzarte (2013)
Spain	Intercepted	NI	Bahillo de la Puebla (2007), Borowski (2007), EPPO (2016a), Fauna Europaea (2016)
Middle East			
Israel	Present	Timber and wood products	Byrinski-Salz (1966), Halperin and Damoiseau (1980), EPPO (2016b)
North America			
USA, New York	Intercepted	Wood boxes	Fisher (1950)
USA, California	Present	Wood, lumber, logs	Ivie (2002) Haack (2006), Hayes and Lundquist (2007), Beiriger (2010)
USA, Florida	Intercepted	Wood products	Woodruff and Fasulo (2006)
South America			
French Guiana	Present	NI	EPPO (2016b)
Asia			
China	Intercepted	Timber imported from Africa	Wei and Guangqin (1994), Zhi-Lin (2003)
Sri-Lanka	Present	Timber	De Silva and Amarasekara (1998)
Oceania			
New Zealand	Intercepted	NI	Archibald and Chalmers (1983)

Present—established and infested. **Intercepted**—identified as introduced in the country (Establishment is doubtful) NI*, No information.

AAYCATAAGAYATYGG-3') and COH6 (5'- TADACTTCDG GRTGDCCAAARAAYCA-3') (Schubart 2009).

Distribution

A literature review was conducted to build a global distribution data where the pest has been present or intercepted before. The primary databases used were Google Scholar, Science Direct, CAB Direct, Web of Science, Zoological Record, and unsystematic exploration using Google search engine. Other than that books and documents

were located at universities and national research center. The keywords used for search were *Heterobostrychus brunneus*, *Ethiopia*, *Cassava pests*, and *tropical wood pests*.

Results and Discussion

Distribution

Information on the geographical distribution of *H. brunneus* is currently limited. The European and Mediterranean Plant Protection




Fig. 2. *H. brunneus* global distribution map (Red, Present; Green, Intercepted). (Source: Authors. Produced by: Quantum GIS Geographic Information System, Version 2.8.2 (Vancouver, British Columbia, Canada). Resolution: 1,000 dpi)




Fig. 3. *H. brunneus* adult (a) Dorsal view, (b) Lateral view and fourth instar larvae (c) Lateral view. (Photos by Aditya Parmar; Collected at Sodo).

Organization (EPP0 2016a,b) and Borowski and Węgrzynowicz (2007) list various countries, namely Cape Verde, French Guiana, Israel, Madagascar, Nigeria, Seychelles, and South Africa. Booth et al. (1990) and Walker (2005) mention Africa as a whole in distribution but do not give specific accounts per country. The geographical distribution of *H. brunneus* provided by ITIS (2016) provides information only at the level of the continent (Africa, Asia, Australia, Europe, and North America) but lacks specific locations. GBIF (2016) lists several countries in Africa and Europe. Fauna Europaea (2016) and Borowski (2007) state the possible presence

(interceptions) of species in Germany, Italy, Spain and United Kingdom. Based on the published information and archives, a list of countries was prepared where *H. brunneus* has been reported previously (Table 1 and Fig. 2). Current literature on stored food product's insects and pests from Ethiopia is devoid of information regarding the presence and infestation of *H. brunneus* (Walker and Boxall 1974, Abate 1988, Tadesse et al. 2006, Hagstrum et al. 2013).

Some of the popular hosts of the insect pest which were listed by Fisher (1950) were bamboo (*Poaceae, Bambusoideae*) in southern




Fig. 4. *H. brunneus* adult: (a) Eye; (b) Thorax lateral view.




Fig. 5. *H. brunneus* adult: (a) Head lateral (or side) view; (b) Pronotum dorsal view.




Fig. 6. Representative holes on cassava chips (a) infested by *H. brunneus* and cassava storage house (b).

Africa; potato from French Guinea and sapwood of various tree species such as African teak (*Baikiaea plurijuga*), Kiaat (*Pterocarpus angolensis*), and Eucalyptus (*Eucalyptus* spp.). *H. brunneus* has been reported to infest stored foodstuffs such as coffee (*Coffea* ssp.) beans, cassava (*M. esculenta*), sweet potato (*Ipomoea batatas*), pulses and oil seeds in tropical regions (Mwanga and Wanyera 1988, Stumpf 1998, Golob et al. 2002, Hill 2002).

Morphology and DNA Barcode

The collected beetle species in cassava storages houses of traders was identified as *Heterobosytrychus brunneus*. Figure 3 provides a pictorial illustration of adult and larvae of the species. Adult beetles have black to dark brownish color and a length of about 7–8 mm and 3–3.5 mm in width. The fourth- to fifth-instar larvae are of pale yellow to whitish color with black mandibles and 9–10 mm long fully elongated, with typical bostrichid profile. Both adults and larvae feed on starchy dried cassava roots.

In the adult beetles, eyes (Fig. 4a) appear marginally detached, upraised and placed on the posterior end of the head (Beiriger 2010). Figure 4b shows the thorax side displaying elytron (dorsal surface) covered lightly with short reclining yellow hairs, are peculiar of *H. brunneus* (Fisher 1950, Walker 2005, Beiriger 2010). The pronotum (Fig. 5b) is much larger than the head (Fig. 5a); the clypeus is flat, and dense yellow hairs are present on the labrum (Fisher 1950). Smooth uneven granules on top of the pronotum (Beiriger 2010), two broadly separated parallel, identical, resembling hooks in front of the pronotum are some of the key features of the species. The sexes of the species are difficult to distinguish based on secondary sexual characteristics. Fisher (1950) states that in the extreme form males have strongly hooked, and a narrowly separated pair of teeth on the anterior margin of the pronotum, whereas female teeth are smaller, straight, and widely separated. However, there are all kinds of variations in between these two extreme forms. To accurately identify the sex of the species, dissecting the specimens and examination of internal genitalia is the only way.

The 658bp long partial sequence of the COI gene of *H. brunneus* was deposited under BOLD Systems database accession number BankIt1918662 CPETH001-16.COI-5P KX232682 (BoldSystems 2016).

Symptoms of Infestation and Management

Cassava tubers in southern Ethiopia are harvested during the dry season (October–February), peeled, chopped into thick slices (2–5 cm) and sun-dried (final moisture content 10–12%). Sun-dried cassava is stored in large quantities until further processing into the composite flour with teff (*Eragrostis tef*) and maize to prepare staple flat bread (Injera). During the survey, 100% of the wholesalers at Sodo and Bale town who were storing cassava for three and more months reported this particular beetle and its larvae as a destructive pest resulting in quantitative and qualitative losses. Turning cassava into powdery dust (Supp. Video S1 and S2 showing adult and larvae respectively feeding on dry cassava), representative entry holes and empty inner spacing were the common symptoms of *H. brunneus* presence. During the cassava (chips and chunks) store inspections at Sodo and Bale, it was observed that the powdery dust and empty inner spacing of cassava were providing a habitat for other grain and flour beetles as well as fungi.

Typical *H. brunneus* bores on dried cassava were about 3–4 mm diameter, illustrated in Figure 6a. Figure 6b shows a common cassava storage system, where cassava chips are filled in polypropylene bags and kept inside or under a roof to protect them from rain.

The common insecticides which were used by traders at the study locations to control *H. brunneus* and other storage insects were Malathion 5% Dust (active ingredient 50g/kg) and Celphos tablets (Aluminium phosphide fumigant insecticide). The presence of *H. brunneus* was reported year round, with not much difference from rainy to the dry season.

Conclusion

Incomplete information is available about the spread and distribution of *H. brunneus* globally. However, the current distribution based on published literature is largely limited to sub-Saharan Africa. Evidence suggests that there is a potential threat that the species may get introduced to and established in other parts of the world through global trade in wood and wood product. Along with its pest status of Wood, *H. brunneus* can be considered as a serious storage pest for dried starchy food products such as cassava. Further studies on damage assessments and sustainable control measures for the species are required in the survey area.

Supplementary data

Supplementary data are available at *Journal of Insect Science* online

Acknowledgments

We thank DAAD (German Academic Exchange Services) and GlobE project Reload (Grant No. 031A247A) (funded by BMBF (Federal Ministry of Education, Germany) and BMZ (Federal Ministry for Economic Cooperation and Development, Germany)). Authors appreciate Prof Rick J. Hodges of NRI (Natural Resource Institute), UK, and Robert Beiriger, Sr Biological Scientist at University of Florida for providing fruitful suggestions on identification and Dr. Helmut Saucke for useful discussion and the provision of the photographic equipment. Further, we thank Dr. Ferdu Azerefege and Dr. Sandip Banerjee of Hawasa University, Hawasa, and Ethiopian Institute of Agricultural Research (EIAR) for co-operation.

References Cited

- Abate, T. 1988. Insect and mite pests of horticultural and miscellaneous plants in Ethiopia. Handbook Nr. 1. Addis Ababa, Ethiopia.
- Alene, D. C., J. Messi, and S. Quilici. 2005. Contribution à la connaissance de la faune d'arthropodes associée à *Ricimodendron heudelotii* Baill. (Euphorbiaceae) au Cameroun. *Fruits* 60: 121–132. (doi: <http://www.fruits-journal.org>)
- Archibald, R. D., and I. Chalmers. 1983. Stored product Coleoptera in New Zealand. *N. Z. Entomol.* 7: 371–397.
- Ashiru, M. 1989. A review of the insect pests and diseases in relation to phenology and distribution of *Triplochiton scleroxylon* K. Schum (Obèche) in West Africa. *Ghana J. Sci.* 29: 371–397.
- Audisio, P., G. Gobbi, G. Liberti, and G. Nardi. 1995. Coleoptera Polyphaga IX (Bostrichoidea, Cleroidea, Lymexyloidea). p. 1–29. In A. Minelli, S. Ruffo, S. La Posta (eds.), Checklist delle specie della fauna italiana, vol. 54. Calderini, Bologna.
- Azmi, M., F. Aboo, and N. Razi. 2011. World distribution of *Heterobosytrychus aequalis* Waterhouse (Coleoptera: Bostrychidae). *J. Entomol.* 8: 497–511.
- Baena, M., and A. J. Zuzarte. 2013. Contribución al estudio de los Bostríquidos de Portugal y actualización del catálogo de la fauna ibérica (Coleoptera, Bostrichidae). *Zool. Baetica.* 25: 25–52.
- Bahillo de la Puebla, P., J. López-Cólon, and M. Baena. 2007. Los Bostrichidae Latreille, 1802 de la fauna de ibero-baleares (Coleoptera). *Heteropterus Rev. Entomol.* 7: 147–227.

- Becker, G. 1980. Wood damage by *Apate monachus* F. and other bostrichid species in West Africa and its (Ivory Coast) prevention. *Mat. Organ.* 15: 245–249.
- Beiriger, R. 2010. *Heterobostrychus hamatipennis* Lesne (Coleoptera: Bostrichidae) new to Florida. *Insecta Mundi*. 0138: 1–5. (<http://journals.fcla.edu>)
- Bellotti, A., and A. van Schoonhoven. 1978. Mite and insect pests of cassava. *Annu. Rev. Entomol.* 23: 39–67.
- Boldsystems. 2016. Sequence page: CPETH001-16. Available at (http://www.boldsystems.org/index.php/MAS_DataRetrieval_OpenSequence?selectedrecordid=7099033).
- Booth, R. G., M. L. Cox, and R. B. Madge. 1990. IIE guides to insects of importance to man. 3. Coleoptera. CAB International Wallingford, UK.
- Borowski, J. 2007. Bostrichidae. *In* L. Löbl, and A. Smetana (eds.), Catalogue of Palaearctic Coleoptera. Volume 4. Elateroidea – Derodontoidea – Bostrichoidea – Lymexyloidea – Cleroidea Cucujoidea. Apollo Books, Stenstrup.
- Borowski, J., and P. Wegrzynowicz. 2007. World catalogue of Bostrichidae Coleoptera. Wydawnictwo Mantis, Olsztyn.
- Brustel, H., and H. P. Aberlenc. 2014. Les Bostrichidae Latreille, 1802 de la faune de France: espèces autochtones, interceptées, introduites ou susceptibles de l'être (Coleoptera). *r.A.R.E.* 23: 65–68.
- BugGuide. 2016. Species *Heterobostrychus brunneus* - Boxwood Borer: Classification. (<http://bugguide.net/node/view/894289>).
- Bytinsky-Salz, H. 1966. Annotated list of the insects and mites introduced into Israel. *Isr. J. Entomol.* 1: 36–37.
- Climate Data Org. 2016. Climate data. (<http://en.climate-data.org/>).
- Cymorek, S. 1970. Eingeschleppte und einheimische Bohr- und Splintholzkafer als Holzschadlinge (Col.; Bostrichidae, Lyctidae): Übersicht zur Lebensweise, über Vorkommen, Einschleppungen, wirtschaftliche Bedeutung, Bekämpfung. *Zeitschrift Fur Angew. Entomol.* 66: 206–224.
- Davies, J. 1960. Coleoptera associated with stored products in Uganda. *East African Agric. J.* 25: 199–201.
- De Silva, N., and H. Amarasekara. 1998. Attack by wood destroying insects on eight commercial timber species, pp. 59–64. *In* H.S. Amarasekera, D.M.S.H.K., Ranasinghe, V. Finlayson (eds.), Proceedings of the Second Annual Forestry Symposium 1996: Management and Sustainable Utilization of Forest Resources. Department of Forestry and Environmental Science, University of Sri Jayewardenepura, Sri Lanka.
- (EPPO) European and Mediterranean Plant Protection Organization. 2016a. (<https://gd.eppo.int/reporting/article-5142>).
- (EPPO) European and Mediterranean Plant Protection Organization. 2016b. EPPO Global Database. *Heterobostrychus brunneus* (HETBBR) (<https://gd.eppo.int/taxon/HETBBR/distribution>).
- Fauna Europaea. 2016. *Heterobostrychus brunneus* (Murray, 1867): Distribution (<http://www.fauna-eu.org>).
- Findlay, W. 1985. Agencies of destruction, part 1, pp. 24–25. *In* W., Findlay (ed.), Preservation of timber in the tropics. Springer Science plus Business Media, Dordrecht.
- Fisher, W. 1950. A revision of the North American species of beetles belonging to the family Bostrichidae. United States Department of Agriculture Miscellaneous Publication, 698, Washington DC.
- Gambetta, A., and E. Orlandi. 1982. Su alcuni insetti reperiti nel legname nei depositi. I° Lictidi e Bostrichidi. *Contrib. Scient.-Prat. Per Una Migliore Conoscenza ed Utilizzazione del Legno.* 30: 1–28.
- (GBIF) Global Biodiversity Information Facility. 2016. (<http://www.gbif.org/species/4427584>).
- Golob, P., G. Farrell, and J. Orchard. 2002. Crop post-harvest: science and technology, Vol. 1. Blackwell Publication Company, UK.
- Gonfa, L. 1996. Climate classifications of Ethiopia. National Meteorological Service Agency, Addis Ababa.
- Haack, R. A. 2006. Exotic bark- and wood-boring Coleoptera in the United States: recent establishments and interceptions. *Can. J. For. Res.* 36: 269–288.
- Hagstrum, D., T. Klejdysz, B. Subramanyam, and J. Nawrot. 2013. Atlas of stored – product Insects and Mites. AACC International, Inc. USA, St. Paul, Minnesota.
- Hagstrum, D., and B. Subramanyam. 2009. Stored-product insect resource. AACC International, Inc. USA, St. Paul, Minnesota.
- Halperin, J., and R. Damoiseau. 1980. The Bostrychid beetles (Coleoptera) of Israel. *Isr. J. Entomol.* 14: 47–53.
- Hayes, J., and J. Lundquist. 2007. The Western bark beetle research group: a unique collaboration with forest health protection, pp. 111–134. *In* Symposium at the 2007 Society of American Foresters Conference. Society of American Foresters, Portland, Oregon.
- Hill, D. 2002. Pest of Stored Foodstuffs and their Control. Kulawer Academic Publisher, Dordrecht, Netherlands.
- Hodges, R. J., J. Meik, and H. Denton. 1985. Infestation of dried cassava (*Manihot esculenta* Crantz) by *Prostephanus truncatus* (Horn) (Coleoptera: Bostrichidae). *J. Stored Prod. Res.* 21: 73–77.
- Howeler, R., N. Ladio, and G. Thomson. 2013. Save and Grow, Cassava: A guide to sustainable production intensification. FAO, Rome, Italy.
- (ITIS) Integrated Taxonomic Information System. 2016. (<http://www.itis.gov>).
- Ivie, M. 2002. Bostrichidae Latreille 1802, pp. 233–244. *In* R.J. Arnett, M. Thomas, P. Skelley, J. Frank (eds.), American beetles polyphaga: scarabaeoidea through curculionoidea. CRC Press, New York.
- Kebede, A., B. Teshome, A. Wondimu, A. Belay, B. Wodajo, and A. Lakew. 2012. Detoxification and consumption of cassava based foods in South West Ethiopia. *Pakistan J. Nutr.* 11: 237–242.
- Lebot, V. 2009. Tropical root and tuber crops: Cassava, sweet potato, yams and aroids. *Crop Prod. Sci. Hortic.* No. 17, CABI Publ. Oxfordshire, UK.
- Lemmens, R. 2012. Baphiopsis. pp. 99–100. *In* R. Lemmens, D. Louppe, and A. Oten-Amoako (eds.), Timbers 2: plant resource of tropical Africa, vol. 7. PROTA Foundation, Wageningen, Netherlands.
- Lesne, P. 1924. Les Coléoptères Bostrychides de l'Afrique tropicale française. *Encycl. Entomol.* 3: 1–301.
- Liu, L., K. Schönitzer, and J. Yang. 2008. A review of the literature on the life history of Bostrichidae (Coleoptera). *Mitteilungen der Münchner Entomologischen Gesellschaft* 98: 91–97.
- Madoffe, S., A. Bakke, and J. Tarimo. 2000. The effect of fire on the diversity and abundance of wood-living beetles in a miombo woodland, Tanzania. *South Afr. For. J.* 187: 51–57.
- Mwanga, R., and N. Wanyera. 1988. Sweet potato growing and research in Uganda, pp. 187–198. *In* Improvement of Sweet Potato (*Ipomoea batatas*) in East Africa, with some references of other tuber and root crops. International Potato Center, Lima, Peru.
- Pringle, J. 1938. Observations on certain wood-boring Coleoptera occurring in South Africa. *Trans. R. Entomol. Soc. London.* 87: 247–270.
- Ratti, E. 2004. Coleoptera Lyctidae e Bostrichidae intercettati nel porto e negli ambienti urbani di Venezia. *Bollettino del Museo Civico di Storia Naturale di Venezia.* 55: 121–125.
- Roberts, H. 1967. A new powder-post beetle, *Xyloperthella guineensis*, together with an annotated Check List of the subfamily Bostrychinae (Col., Fam. Bostrichidae) from Nigeria. *J. Nat. Hist.* 2: 85–104.
- Robinson, W. H. 2005. Urban insects and arachnids - a handbook of urban entomology. Cambridge University Press, Cambridge, UK.
- Schabel, H. G. 2006. Forest entomology in East Africa: forest insects of Tanzania. Springer, Dordrecht, Netherlands.
- Schubart, C. D. 2009. Mitochondrial DNA and decapod phylogenies: The importance of pseudogenes and primer optimization, pp. 47–65. *In* J. W. Martin, K. A. Crandall and D. L. Felder (eds), Crustacean issues 18 – decapod crustacean phylogenetics. Taylor and Francis/CRC Press Boca Raton, Florida
- Selander, J., and G. Pearce. 1984. Seasonal life history and economic importance of the major timber beetles of the Zambian teak forest, pp. 287–301. *In* First International Conference on the Teak Forests of Southern Africa. Livingstone, Zambia.
- Stumpf, E. 1998. Post-harvest loss due to pests in dried cassava chips and comparative methods for its assessment: a case study on small-scale farm households in Ghana. Ph.D. thesis, Humboldt-Universität, Berlin. (<http://www.fao.org/wairdocs/>)
- Tadesse, A., A. Ayalew, E. Getu, and T. Tefera. 2006. Review of research on post-harvest pests. pp. 475–563. *In* A. A., Tadesse (ed.), Increasing crop

- production through improved plant protection Vol. 2. Plant protection society of Ethiopia, Ethiopia.
- Vrydagh, J. 1955a. Contribution à l'étude des Bostrychidae n.4. Collection du Musée zoologique de "Humboldt Universitat" à Berlin. Bull. Soc. R. Belge D'Entomol. 31: 1–16.
- Vrydagh, J. 1955b. Contribution à l'étude des Bostrychidae Coleoptera Teredilia 3. - Les Bostrychidae du Mozambique. Bull. Soc. R. Belge D'Entomol. 31: 1–23.
- Vrydagh, J. 1960. Contribution à l'étude des Bostrychidae. 23. — Collection de la Section Zoologique du Musée National Hongrois à Budapest. Bull. Inst. R. des Sci. Nat. Belgique. 36: 1–32.
- Wagner, M. R., J. R. Cobbinah, and P. P. Bosu. 2008. Forest entomology in West Tropical Africa: forests insects of Ghana. Springer Science & Business Media. Dordrecht, Netherlands.
- Walker, D. J., and R. A. Boxall. 1974. An annotated list of the insects associated with stored products in Ethiopia, including notes on mites found in Harar Province. East Afr. Agric. Forest. J. 39: 330–335.
- Walker, K. 2005. Auger beetle (*Heterobostrychus brunneus*) Updated on 10/21/2011 9:42:24 AM Available online: PaDIL – (<http://www.padil.gov.au>).
- Wasonga, V., P. Malaki, R. Mwakodi, and K. Matheka. 2015. Fifth biodiversity assessment of Ol Ari Nyiro, Laikipia nature conservancy, northern, Kenya (Amphibians, Reptiles, Birds, Invertebrates and Plants). National museums of Kenya, Nairobi.
- Wie, X., and L. Guangqin. 1994. The catch and disinfection of Bostrychidae. Acta Agric. Univ. Jiangxiensis. 4: 362–366.
- Woodruff, R., and T. Fasulo. 2006. An Oriental Wood Borer, *Heterobostrychus aequalis* (Waterhouse) (Insecta, Coleoptera, Bostrychidae). University of Florida, IFAS Extension, 3 pp.
- Wylie, F. R., M. Griffiths, and J. King. 2008. Development of hazard site surveillance programs for forest invasive species: a case study from Brisbane, Australia. Aust. For. 71: 229–235.
- Zhi-Lin, C. 2003. Classification of the Bostrichidae pests intercepted in China on timber imported from Africa. Entomol. Knowl. 40: 154–159.